

Surgeon's Neck Posture during Spine Surgeries: "The Unrecognised Potential Occupational Hazard"

Abstract

Study Design: Observational study. **Purpose:** The purpose of this study is to analyze the surgeon's neck postures while performing lumbar spinal surgeries. **Overview of Literature:** Lumbar spinal surgeries are on rising trend, and with increase in number of procedures, the average time spent by a spine surgeon performing surgical procedures is also increasing. The effect of operating posture on the surgeon's neck is largely unknown. From the studies conducted on usage of smartphones, abnormal neck postures, especially the forward head posture (FHP), were found to adversely affect the cervical spine of individuals. The present study analyzes the neck position of spine surgeons during lumbar spine surgeries. **Methodology:** Sixty video recordings (25 open transforaminal lumbar interbody fusions [TLIFs] and 35 lumbar decompression [LD] procedures – 15 with headlight and 20 with operating microscope) of surgeries performed by three spine surgeons of different heights were analyzed. Running videos of the surgeries were recorded concentrating on the surgeons with reflective markers taped to their surface landmarks corresponding to C7 spinous process, tragus of the ear, and outer canthus of the eye. Video recordings were standardized by a fixed video recorder in the same operating theater. Snapshots from the video were obtained whenever the surgeon changes the position. Head flexion angle (HFA), neck flexion angle (NFA), and cervical angle (CA) were measured and analyzed. **Results:** During TLIF, HFA and NFA were significantly higher during the phases of decompression and fusion ($P < 0.05$). The average CA of all surgeons was lower, thereby adversely affecting the cervical spine ($20.15^\circ \pm 5.05^\circ$). During LD, CA showed significant difference between usage of microscope and headlight ($P < 0.001$). **Conclusion:** Surgeon's FHP is frequently caused by a compromise between the need to perform surgery with hands, without elevating the arms, and simultaneous control of gaze at surgical field. The usage of microscope was found to reduce the stress on neck while performing surgery.

Keywords: Ergonomics, forward head posture, postural analysis, spine surgeon, surgeon's neck

**J Naresh-Babu,
Viswanadha
Arun-Kumar,
D G S Raju**

*Mallika Spine Centre, Guntur,
Andhra Pradesh, India*

Introduction

Lumbar spine surgeries are on a rising trend in the past few decades, the reasons being many from advancement of instrumentation technologies to experience of the surgeons. As a reason, the average time spent by a spine surgeon performing surgical procedures is also increasing. Since intraoperative position of one's own neck is not regularly assessed or observed by the spine surgeons, the effect of operating posture on the surgeon's neck is largely unknown.

Good posture is defined by the Posture Committee of the American Academy of Orthopedic Surgeons as "the state of muscular and skeletal balance which protects the supporting structures of the body against injury or progressive

deformity, irrespective of the position (erect, lying, squatting, or stooping) in which these structures are working or resting".¹ Correct upright posture is defined as when ears are aligned with the shoulders in the same line, leading to least strain on the back when in standing position.² To maintain such a good posture, one must be always aware of the position in which he/she is constantly working and correct it on regular basis.

Postural neck pain can be caused by several factors, including sleeping with head elevated too high, overuse of computers or cell phones, and lack of back muscle strength.³ The common neck-related syndromes are straight neck syndrome and forward head posture (FHP). Loss of natural cervical curve is termed as straight neck syndrome, which is mainly asymptomatic but when neglected can lead

Address for correspondence:

*Dr. Viswanadha Arun-Kumar,
Mallika Spine Centre,
12-12-30, Old Club Road,
Kothapet, Guntur - 522 001,
Andhra Pradesh, India.
E-mail: drarunmsortho@gmail.
com*

Access this article online

Website: www.ijoonline.com

DOI:
10.4103/ortho.IJOrtho_677_18

Quick Response Code:



How to cite this article: Naresh-Babu J, Arun-Kumar V, Raju DG. Surgeon's neck posture during spine surgeries: "The unrecognised potential occupational hazard". Indian J Orthop 2019;53:758-62.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

to pain-associated FHP. According to Hansraj,⁴ weight load on the spine dramatically increases when there is an increase in neck flexion ranging from 10 lbs at 0° to 60 lbs at 60°. It is around 10–12 lbs at neutral position and 27, 40, 49, and 60 lbs at 15°, 30°, 45°, and 60°, respectively. According to Szeto *et al.*,⁵ when the head is in forward flexion posture for a longer time period, it results in loss of cervical lordosis and potential increase in thoracic kyphosis. FHP combined with increased biodynamic stress of the spine leads to musculoskeletal problems, such as neck pain and headache.⁶ There have been reports where temporomandibular dysfunction is also associated with FHP.⁷ Several studies have also come up examining the influence of head weight on flexion head posture.^{8,9}

Postural analysis has been extensively studied over the last decade, with many development methods gaining importance. The latest reliable developmental methods used advanced technological systems such as computerized photographic systems¹⁰ and X-ray scannograms.^{11,12} Methods to evaluate spinal posture have been categorized basically into four groups:

1. Radiography¹³ – Reliable and gold standard but involves radiation hazards
2. Three dimensional motion analysis¹⁴ – Reliable but requires costly equipment
3. Video raster stereography analysis¹⁵ – Reliable but did not pass validity studies
4. Photographic posture analysis¹⁶ – Basic objective observational measurement method using anatomical landmarks.

Reliability and clinical use of the photographic analysis have been described in the literature based on its accuracy.^{10,17} According to van Niekerk,¹⁸ photographs are reliable indicators of the spine when compared to radiographs using LODOX system.

The purpose of the study is to analyze the effect of FHP on the neck of individuals performing spinal surgeries.

Methodology

An observational study of 60 videos performed by three spine surgeons (S1, S2, and S3) was conducted at our institute. Heights of the surgeons were calculated using photogrammetry and Photographic Posture Analysis Method by single research analyst. Preoperatively, reflective markers were taped on the side of the surgeons [Figure 1], in the following surface landmarks:

1. C7 spinous process
2. Tragus of the ear
3. Outer canthus of the eye.

Reflective markers used were Styrofoam balls of approximately 20-mm diameter. Running video of the surgery was recorded using NIKON Powershot D3100 placed on a Manfrotto tripod at an angle perpendicular to the operative field concentrating on the performing surgeon



Figure 1: Reflective markers taped on side of the surgeon

primarily. The distance between the field and the camera was set at 2.5 metres. Height of the tripod was adjusted so as to center the head and neck of the surgeon in the field. A precalibrated board was placed in the field of view to allow referencing vertical and horizontal axes from the photograph. Postoperatively, all the videos involving three surgeons were analyzed by single research analyst. The total number of operative videos analyzed was 60 which included 35 lumbar decompression (LD) procedures and 25 transforaminal lumbar interbody fusions (TLIFs). Duration of the whole surgery of TLIF was divided into different phases as exposure, fixation, decompression, fusion, and closure. Videos of LD performed were grouped into Group H – surgery performed with usage of headlight and Group M – surgery performed under microscope. Snapshots of the video were taken whenever the surgeon changes his/her position, and using Surgimap (Spine Software, version 2.2.9.9.4, New York, NY, USA), all images were calibrated. The angles evaluated in the study are described below [Figure 2a-c]:

1. Head flexion angle (HFA) is the angle between a line connecting C7 to tragus of the ear and tragus to outer canthus¹⁹
2. Neck flexion angle (NFA) is defined as the angle subtended between vector pointing from C7 to tragus (which corresponds to occipital-cervical joint) and a global vertical line²⁰
3. Cervical angle (CA) has been one of the reliable indicators to assess FHP and is defined as the angle formed at the intersection of the horizontal line through the spinous process and tragus of the ear.²¹

Results

Analysis of posture during transforaminal lumbar interbody fusions

The average duration of surgery for single-level TLIF is 110.63 min (range 71.77–148.36 min). The average distance between surgeon's center of ear and center

of shoulder during the entire surgery is 135.5 mm (range 97.82–190.81 mm) [Table 1].

HFA of all surgeons was observed to be near-normal (S1 – 116.5° ± 6.4°, S2 – 122° ± 5.9°, S3 – 130.6° ± 5.4°) during exposure, abnormally increased (S1 – 145° ± 3.4°, S2 – 143.8° ± 2.9°, S3 – 149.7° ± 4.1°) during decompression, and reduced back to near-normal range (S1 – 120.3° ± 3.0°, S2 – 126.6° ± 2.5°, S3 – 132.3° ± 3.4°) during wound closure.

NFA of the surgeons was also observed to follow a similar trend as of HFA, which showed to be abnormal during decompression (S1 – 76.7 ± 4.2, S2 – 80.2 ± 4.1, and S3 – 98.7 ± 4.5) and fusion (S1 – 68.4 ± 5.3, S2 – 79.7 ± 3.3, S3 – 93.6 ± 4.2).

Decompression and fusion were observed to be the most stressful phases affecting surgeon's neck during TLIF. HFA and NFA were significantly higher during the phases of decompression and fusion when compared with exposure and closure (*P* < 0.05). The average CA of all surgeons was significantly lower in all phases of the entire surgery, thereby adversely affecting the cervical spine (20.15° ± 5.05°).

Analysis of posture during decompression

HFA and NFA did not alter much in the headlight and microscope groups, while CA showed significant difference (*P* < 0.001). With the usage of microscope, CA was observed to be near-normal for the entire duration [Table 2].

Discussion

Neck-related syndromes are rapidly increasing as the lifestyle of majority of the population involves looking down using smartphones and longer duration of working hours in front of a computer. FHP being a trending condition affecting the younger generation due to usage of technologies such as smartphones, this can also affect the spine surgeons to a particular level. The number of spine surgeries performed has increased markedly in the past decade due to many innovations in the specialty field.^{22,23}




Figure 2: Snapshot pictures showing head flexion angle (a), neck flexion angle (b), and cervical angle (c) of operating surgeons

Table 1: Results showing head flexion angle, neck flexion angle, and cervical angle of operating surgeons during different phases of lumbar fixation and fusion

	HFA		NFA		CA		Distance	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Exposure								
Surgeon 1	116.5	6.4334	55	4.02768	28.6	2.319	106.93	3.459
Surgeon 2	122	5.9066	67	3.944	20.5	1.7795	138.08	1.5604
Surgeon 3	130.6	5.4405	73	4.2018	17.3	1.6329	177.16	2.1498
FIXATION								
Surgeon 1	131	3.8005	60.7	4.5472	25.1	3.2472	110.22	2.8312
Surgeon 2	139.8	2.7406	73.2	3.2591	19.7	2.3118	151.03	1.8506
Surgeon 3	141.3	3.4657	87.6	3.4058	14.2	2.1421	179.05	3.2903
Decompression								
Surgeon 1	145	3.496	76.7	4.2176	21.2	2.8596	159.67	2.8627
Surgeon 2	143.8	2.9363	80.2	4.1311	18.2	2.6583	164.17	3.2964
Surgeon 3	149.7	4.139	98.7	4.5227	12.3	2.5841	190.81	2.56476
FUSION								
Surgeon 1	135.9	3.6651	68.4	5.3374	23.8	4.1311	136.53	2.808
Surgeon 2	142.2	4.1311	79.7	3.3349	18	2.7487	157.91	2.9898
Surgeon 3	150.8	4.8716	93.6	4.2216	12.7	1.7029	187.41	2.3278
Closure								
Surgeon 1	120.3	3.093	57.36	3.4719	28	2.9059	97.82	4.0746
Surgeon 2	126.6	2.5033	67.3	2.6687	24	3.055	137.34	3.4296
Surgeon 3	132.3	3.4657	77.7	3.4334	18.7	3.3747	157.56	3.9699

Table 2: Results of lumbar discectomy performed with headlight and microscope

	HFA		NFA		CA		Distance	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Head light								
Surgeon 1	111.4	2.6331	55.5	3.3747	22	3.1269	110.96	3.3118
Surgeon 2	120.5	2.9533	58.9	4.1486	17.1	3.6651	127.1	2.9137
Surgeon 3	138	2.708	73.7	3.2676	14.4	3.2041	140.09	3.4296
Microscope								
Surgeon 1	108.5	2.9907	57.6	3.7771	38.1	3.0713	111.35	3.1502
Surgeon 2	116.1	3.5418	59.7	3.1287	36.4	3.1692	120.45	3.0297
Surgeon 3	120.9	3.9258	65.8	2.8982	31.1	2.8067	130.66	2.3527

If prolonged work is performed with neck in flexion, pain associated with fatigue is the common symptom. Main load on the neck musculature is transferred to surrounding ligaments and capsules of cervical facet joints when head is flexed forward to extreme of its range of motion. Such postures can lead to pain within 15 min, and there is a usual tendency to normalize the posture within 15–60 min due to intense pain. Pressure exerted by head and neck on the disc between the seventh cervical vertebra and the first thoracic vertebra is increased by 3.6 times when the cervical spine is forward flexed maximally.²⁴

Dangers of FHP can be described as a “DOMINO EFFECT” since one effect sets of a chain of similar events, leading to a cumulative result, i.e.,

1. Head moves forward, thereby shifting the center of gravity
2. As a primary compensatory mechanism, upper body drifts backward
3. As a secondary compensation to upper body drift, both hips tilt forward.

Hence, FHP can not only cause neck pain but can also be a root cause of mid/lower back pain.

The relation of head to that of the neck can be determined by the following parameters: NFA and CA. According to Yoo,²⁵ NFA at $38.5^\circ \pm 6.02^\circ$ increases muscle activity of the upper trapezius and splenius capitis muscles, when compared to cervical neutral position. In a series by Lee,²⁶ muscle fatigue levels of the right and left upper trapezius was highest at 50° and lowest at 30° . In our study, NFA was above 50° in all instances proving that muscle fatigue levels of trapezius is highest during both TLIF and LD.

Many studies stated that normal individuals have mean CA ranging between 40° and 55° .^{27,28} In the series by Ruivo *et al.*,²¹ cases with FHP associated with neck pain had a CA $<50^\circ$. Brink *et al.*²⁹ found smaller CA ($<40^\circ$) to be the reason behind upper quadrant pain. In the present study, we observed the CA remained $<25^\circ$ for all the surgeons for the entire duration of the surgery, irrespective of the phase. Smaller the CA indicates more the FHP.

According to Kessel,³⁰ for every inch of FHP, there is an increase in weight of head on the spine by additional

10 pounds. In our study, the average FHP of operating surgeon has been 5.3 inch which amounts to >50 pounds of head weight on the cervical spine. Normally, the weight load of head on spine is 10–12 lbs where the neck flexion is at neutral position and center of the ear lies in the same line that of center of shoulder blades. In the series by Hansraj conducted on bone models, when the neck flexion increases to 60° , load becomes 60 lbs and from here on load could not be calculated since the modules were termed becoming unstable in higher flexion degrees. In the present study, load of head on the spine of surgeons has been consistently above 60 lbs during all the phases of surgery based on the findings of NFA.

Surgeon's FHP is frequently caused by a compromise between the need to perform surgery with hands, without elevating the arms and simultaneous control of gaze at surgical field. The phases showing maximum effect of FHP on the neck of the surgeon are decompression and interbody fusion, whereas exposure and closure of the wound are the least affected phases in the surgery. On an average, if a highly experienced surgeon (>10 years) performs five single-level interbody fusions per week, he/she is expected to be in the position of FHP for about 311.00 hours/year. When the same number of cases if being done by moderately experience surgeon (5 years), they are expected to be in the same abnormal position for about 484.29 hours/year and least experienced (<2 years) time duration would be 642.89 hours/year, which is highly significant when compared to that of the highly experienced spine surgeons. Furthermore, the usage of microscope was found to be beneficial by avoiding the abnormal neck posture angles when compared to the usage of headlight.

Consistent flexion of the spine has been associated with increased cervical compressive loading and reprobate response in the cervical connective tissues.^{31,32} The angle of neck flexion in a surgeon that is observed during performing spine surgery is causing a severe load on the surgeon's spine. To avoid excessive consistent pressure on the neck, we suggest changing the posture regularly after each phase for at least 10 s and further recommend spine surgeons to actively perform neck strengthening exercises on a daily basis, which will decrease the effect of FHP on surgeon's neck in the years to come.

Conclusion

When the neck stays in such an abnormal position on a daily basis, there is a huge pressure on the surgeon's neck making it highly vulnerable for early degeneration. Usage of microscope was found to reduce the stress on neck while performing surgery. The data of this study creates an awareness among spine surgeons about occupational hazards and need for regular neck strengthening exercises.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

References

- Grimmer-Somers K, Milanese S, Louw Q. Measurement of cervical posture in the sagittal plane. *J Manipulative Physiol Ther* 2008;31:509-17.
- Straus EW. The Upright Posture. In: Natanson M. (eds) *Essays in Phenomenology*. Springer, Dordrecht. 1966. doi: doi.org/10.1007/978-94-017-5403-3_10.
- Edmondston SJ, Wallumrød ME, Macleód F, Kvamme LS, Joebges S, Brabham GC. Reliability of isometric muscle endurance tests in subjects with postural neck pain. *J Manipulative Physiol Ther* 2008;31:348-54.
- Hansraj KK. Assessment of stresses in the cervical spine caused by posture and position of the head. *Surg Technol Int* 2014;25:277-9.
- Szeto GP, Straker L, Raine S. A field comparison of neck and shoulder postures in symptomatic and asymptomatic office workers. *Appl Ergon* 2002;33:75-84.
- Fernández-de-las-Peñas C, Alonso-Blanco C, Cuadrado ML, Pareja JA. Forward head posture and neck mobility in chronic tension-type headache: A blinded, controlled study. *Cephalalgia* 2006;26:314-9.
- Hanten WP, Lucio RM, Russell JL, Brunt D. Assessment of total head excursion and resting head posture. *Arch Phys Med Rehabil* 1991;72:877-80.
- Burgess-Limerick R, Plooy A, Ankrum DR. The effect of imposed and self-selected computer monitor height on posture and gaze angle. *Clin Biomech (Bristol, Avon)* 1998;13:584-92.
- Sarig-Bahat H. Evidence for exercise therapy in mechanical neck disorders. *Man Ther* 2003;8:10-20.
- Ferreira EA, Duarte M, Maldonado EP, Burke TN, Marques AP. Postural assessment software (PAS/SAPO): Validation and reliability. *Clinics (Sao Paulo)* 2010;65:675-81.
- Wunderlich M, Rütger T, Essfeld D, Erren TC, Piekarski C, Leyk D, *et al.* A new approach to assess movements and isometric postures of spine and trunk at the workplace. *Eur Spine J* 2011;20:1393-402.
- Brink Y, Louw Q, Grimmer-Somers K. The quality of evidence of psychometric properties of three-dimensional spinal posture-measuring instruments. *BMC Musculoskelet Disord* 2011;12:93.
- Vedantam R, Lenke LG, Keeney JA, Bridwell KH. Comparison of standing sagittal spinal alignment in asymptomatic adolescents and adults. *Spine (Phila Pa 1976)* 1998;23:211-5.
- Uritani D. Reliability of upper quadrant posture analysis using an ultrasound-based three-dimensional motion analyzer. *J Phys Ther Sci* 2013;25:1181-4.
- McEvoy MP, Grimmer K. Reliability of upright posture measurements in primary school children. *BMC Musculoskelet Disord* 2005;6:35.
- Nam SH, Son SM, Kwon JW, Lee NK. The intra-and inter-rater reliabilities of the forward head posture assessment of normal healthy subjects. *J Phys Ther Sci* 2013;25:737-9.
- Fortin C, Feldman DE, Cheriet F, Labelle H. Clinical methods for quantifying body segment posture: A literature review. *Disabil Rehabil* 2011;33:367-83.
- van Niekerk SM, Louw Q, Vaughan C, Grimmer-Somers K, Schreve K. Photographic measurement of upper-body sitting posture of high school students: A reliability and validity study. *BMC Musculoskelet Disord* 2008;9:113.
- Burgess-Limerick R. The influence of computer monitor height on head and neck posture. *Int J Ind Ergon* 1999;23:171-9.
- Young JG, Trudeau M, Odell D, Marinelli K, Dennerlein JT. Touch-screen tablet user configurations and case-supported tilt affect head and neck flexion angles. *Work* 2012;41:81-91.
- Ruivo RM, Pezarat-Correia P, Carita AI. Cervical and shoulder postural assessment of adolescents between 15 and 17 years old and association with upper quadrant pain. *Braz J Phys Ther* 2014;18:364-71.
- Rajae SS, Bae HW, Kanim LE, Delamarter RB. Spinal fusion in the United States: Analysis of trends from 1998 to 2008. *Spine (Phila Pa 1976)* 2012;37:67-76.
- Davis H. Increasing rates of cervical and lumbar spine surgery in the United States, 1979-1990. *Spine (Phila Pa 1976)* 1994;19:1117-23.
- Stellman JM, Editor. *Encyclopaedia of occupational health and safety*. International Labor Organisation, Geneva; 1998. Available from: <http://iloencyclopaedia.org/component/k2/17-6-musculoskeletal-system/neck>. [Last accessed on 2019 Jul 15].
- Yoo CU. Electromyographic Activity of the Neck and Shoulder Muscles While Watching a DMB Phone with the Neck Flexed. Yonsei University, Dissertation of Master's Degree; 2008.
- Lee S, Lee D, Park J. Effect of the cervical flexion angle during smart phone use on muscle fatigue of the cervical erector spinae and upper trapezius. *J Phys Ther Sci* 2015;27:1847-9.
- Lee MY, Lee HY, Yong MS. Characteristics of cervical position sense in subjects with forward head posture. *J Phys Ther Sci* 2014;26:1741-3.
- Lau HM, Chiu TT, Lam TH. Measurement of craniovertebral angle with electronic head posture instrument: Criterion validity. *J Rehabil Res Dev* 2010;47:911-8.
- Brink Y, Crous LC, Louw QA, Grimmer-Somers K, Schreve K. The association between postural alignment and psychosocial factors to upper quadrant pain in high school students: A prospective study. *Man Ther* 2009;14:647-53.
- Kessel L. The physiology of the joints. Volume 3. The trunk and the vertebral column. I. A. Kapandji, Paris. Second edition. 1974. Edinburgh: Churchill Livingstone. *Br J Surg* 1975;62:419.
- Harms-Ringdahl K, Ekholm J, Schüldt K, Németh G, Arborelius UP. Load moments and myoelectric activity when the cervical spine is held in full flexion and extension. *Ergonomics* 1986;29:1539-52.
- Twomey L, Taylor J. Flexion creep deformation and hysteresis in the lumbar vertebral column. *Spine (Phila Pa 1976)* 1982;7:116-22.

Copyright of Indian Journal of Orthopaedics is the property of Wolters Kluwer India Pvt Ltd and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.